

In this issue

Research Article

[Open Access](#) [Research Article](#) PTZAID:AHCRR-4-119

[Myeloproliferative Neoplasms: Fatigue, depression & hemoglobi](#)

Published On: July 02, 2019 | Pages: 014 - 019

Author(s): Marina Stal*

Objective: To evaluate existing relationships and differences in fatigue, depression, and hemoglobin value in individuals with Myeloproliferative Neoplasms. Methods: This study utilized the Fatigue Severity Scale, Beck Depression Inventory-II, and self-report of MPN diagnosis and latest hemoglobin value to assess the dependent variables. To promote optimal opportuni ...

[Abstract View](#) | [Full Article View](#) | DOI: [10.17352/ahcrr.000019](https://doi.org/10.17352/ahcrr.000019)

[Open Access](#) [Research Article](#) PTZAID:AHCRR-4-116

[Synergy of blood grouping and body sweating](#)

Published On: April 09, 2019 | Pages: 006 - 007

Author(s): Muhammad Imran Qadir and Muhammad Asad*

Objective of the present study was to correlate blood grouping with “sweating in body”. ABO typing is the test to determine your blood group. In this test, you had to mix your blood with antibodies which were against the type of blood A and blood B. Blood typing is very important during pregnancy. When you had done the blood typing you have to see either Rh factor is ...

[Abstract View](#) | [Full Article View](#) | DOI: [10.17352/ahcrr.000016](https://doi.org/10.17352/ahcrr.000016)

[Open Access](#) [Research Article](#) PTZAID:AHCRR-4-114

[Outcome of blood group system on selfie loving](#)

Published On: February 07, 2019 | Pages: 001 - 002

Author(s): Muhammad Imran Qadir and Muhammad Hamad Saif*

Objective of the present study was co-relating blood grouping with selfie lover. A questionnaire was prepared about Selfie lover. Total numbers of students were 169 students in this research and the average age of students was 18-22 years. All the students give answer according to their blood group. Results completed according to project. We took blood sample of every ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/ahcrr.000014

Case Report

[Open Access](#) | [Case Report](#) | PTZAID:AHCCR-4-120

[A case of severe dermatitis in a patient with Polycythemia Vera during cytoreductive therapy](#)

Published On: July 20, 2019 | Pages: 020 - 021

Author(s): Accurso V, Santoro M*, Caputo V, Fiorella S, Casimiro P, Sardo M, Marino C, Sucato G and Siragusa S

Polycythemia Vera (PV) is a Philadelphia-negative chronic myeloproliferative neoplasm (MPN) mainly characterized by erythrocytosis. In this report we describe a case of severe cutaneous toxicity in patients with PV treated with hydroxyurea. A 72-year-old woman diagnosed with PV with V617F mutation of JAK2 performed more than 10 years before and treated with hydroxy ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/ahcrr.000020

[Open Access](#) | [Case Report](#) | PTZAID:AHCCR-4-118

[Dissemine intravascular coagulation may be the presenting feature for chronic myelomonocytic leukemia: Special case report](#)

Published On: July 01, 2019 | Pages: 011 - 013

Author(s): Itir Sirinoglu Demiriz*

Chronic myelomonocytic leukemia (CMML) is a malignant myeloid stem cell disease accompanied by dysplasia in the context of myeloproliferative disease. Peripheral cytopenias (mainly anemia and thrombocytopenia) and hepatosplenomegaly are common findings. ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/ahcrr.000018

[Chronic demyelinating polyneuropathy associated with anti-ganglioside GQ1b antibodies in peripheral T-cell lymphoma](#)

Published On: May 23, 2019 | Pages: 008 - 010

Author(s): Salem Bahashwan*, Carolyne Croizier, Xavier Moisset, Romain Guièze, Bertrand Evrard, Olivier Tournilhac, Jacques-Olivier Bay and Richard Lemal

Malignant lymphoma can cause peripheral neuropathy through various mechanisms. We report the case of 67-year-old man with chronic motor and sensory axonal demyelinating polyneuropathy associated with anti-ganglioside GQ1b antibodies in serum and cerebrospinal fluid (CSF) as an initial presentation of peripheral T-cell lymphoma, not otherwise specified. The patient was ...

[Abstract View](#) | [Full Article View](#) | DOI: [10.17352/ahcrr.000017](https://doi.org/10.17352/ahcrr.000017)

[Recurrent thrombosis in a multiple myeloma patient with low risk for thrombosis despite being on thromboprophylaxis](#)

Published On: February 19, 2019 | Pages: 003 - 005

Author(s): Chow Siew Kian* and Fu Mei Sian

It is a well-known fact that multiple myeloma (MM) patients have an increased risk for thromboembolic events involving both the venous and arterial territories. The etiologies can be multifactorial, for example, activation of procoagulant factors, acquired protein C resistance, MM-specified and treatment-related factors. The risk of venous thrombosis is high when pati ...

[Abstract View](#) | [Full Article View](#) | DOI: [10.17352/ahcrr.000015](https://doi.org/10.17352/ahcrr.000015)