

In this issue

Case Report

[Open Access](#) [Case Report](#) PTZAID:GJMCCR-5-160

[The Anemia that Halted Treatment: A case of Complicated Hyperlipidemia](#)

Published On: March 22, 2018 | Pages: 010 - 013

Author(s): Mark M Mokhtar*, Catherine E Lempke and Ruth Agwuna

Familial hyperlipidemia is defined by abnormal levels of the following: low density lipoprotein cholesterol (LDL-C); high density lipoprotein cholesterol (HDL); triglycerides (TAG); or any combination of the three. ...

[Abstract View](#) [Full Article View](#) [DOI: 10.17352/2455-5282.000060](#)

[Open Access](#) [Case Report](#) PTZAID:GJMCCR-5-159

[The Heart That Wasn't Attacked: A Case of Transverse Myelitis](#)

Published On: March 16, 2018 | Pages: 007 - 009

Author(s): Mark M Mokhtar*, Ahmed Ibrahim and Catherine E Lempke

Transverse myelitis is an autoimmune demyelination disease of the spinal cord that can present with a myriad of symptoms ranging from altered sensations, weakness, loss of bowel control and so forth. There are approximately 1400 new cases reported each year in the United States, but this is likely an underestimation, without a gender bias or familial predisposition a ...

[Abstract View](#) [Full Article View](#) [DOI: 10.17352/2455-5282.000059](#)

[Open Access](#) [Case Report](#) PTZAID:GJMCCR-5-158

[Spontaneous Rupture of the Kidney Due to Acute Lymphoblastic Leukemia: The First Episode and a Glance on the Literature](#)

Published On: February 05, 2018 | Pages: 003 - 006

Author(s): Sami Mahjoub Taha*, Awad Al Kareem Abd El Geyoom, Mohammed El Imam M Ahmed, Nada Osman and Musab Mahjoub Taha

Spontaneous rupture of the kidney consists of the atraumatic disruption of the renal parenchyma, followed by hemorrhage or leakage of urine into the perinephric space. ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-5282.000058

[Open Access](#) | [Case Report](#) | PTZAID:GJMCCR-5-157

[A giant abdominal well differentiated Liposarcoma: A rare etiology of abdominal mass](#)

Published On: January 31, 2018 | Pages: 001 - 002

Author(s): Soukaina EL Yazal*, Abdelouahed Louzi, Fatima Z Lairani, Soukaina Jiddi, Imane Haraki, Sofi a Oubaha, Zouhour Samlani Sebbane and Khadija Krati

Liposarcoma is the most common retroperitoneal sarcoma and represents more than 15% of all sarcomas. Well differentiated liposarcoma is usually located in the extremities. We report the case of a patient with a giant retroperitoneal liposarcoma revealed by an abdominal mass. ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-5282.000057