

In this issue

Research Article

[Open Access](#) [Research Article](#) PTZAID:IJICR-6-127

[Selecting patients for treatment with immunotherapy: Our experience in a resource-deprived economy](#)

Published On: October 27, 2020 | Pages: 025 - 027

Author(s): Gabriel Chianakwana*, Amobi Egwuonwu, Christian Okoli, Eric Ihekwoaba, Dickson Emegoakor, Henry Nzeako, Evaristus Afiadigwe and Dan Anyiam

Background: Majority of the solid cancer patients seen in our Centers come with advanced diseases. Some of the cancers are locally advanced, while others are metastatic. Most of these patients who present late belong to the poverty-stricken group in our community. Only few of them can afford costly drugs for their treatment. The government Health Insurance Scheme does ...

[Abstract View](#) [Full Article View](#) [DOI: 10.17352/2455-8591.000027](#)

[Open Access](#) [Research Article](#) PTZAID:IJICR-6-122

[Tadalafil inhibits elevated glutamic oxaloacetic transaminase during alcohol aflatoxin induced hepatocellular carcinoma in rats](#)

Published On: February 25, 2020 | Pages: 010 - 013

Author(s): Divya Rawat and Raj Kumar Koiri*

Hepatocellular carcinoma is a type of primary liver cancer and dietary exposure to aflatoxins is one of major causative factor for the development of HCC. Present study was planned to assess the role of PDE5 inhibitor tadalafil on glutamic oxaloacetic transaminase during aflatoxin induced HCC. Rats of control group received normal food and water ad libitum. Alcohol an ...

[Abstract View](#) [Full Article View](#) [DOI: 10.17352/2455-8591.000022](#)

[Open Access](#) [Research Article](#) PTZAID:IJICR-6-121

[Evaluation of breast cancer regarding molecular and immunochemical markers](#)

Published On: February 20, 2020 | Pages: 001 - 009

Author(s): Sally Ahmad Sabra, Abir Adel Saad, Nadia Ahmad Abd El Moneim, Mahmoud Abd El Atty Hemida, Nermine Moussa and Medhat Haroun*

Early-stage cancer recognition could improve awareness and treatment strategies. Twenty four breast cancer affected individuals had their DNA isolated from malignant tissues and from blood. DNA was then amplified by RAPD with six different ten-mer primers. RAPD-PCR yields were electrophoresed on a 1 .5% agarose gel and visualized using ethidium-bromide staining. Only ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-8591.000021

Review Article

[Open Access](#) | [Review Article](#) | PTZAID:IJICR-6-128

[The Biomarkers of Cd4+ T Regulatory Cells Associated with Tumour Immune Escape](#)

Published On: October 28, 2020 | Pages: 028 - 034

Author(s): Nyaribari MC*

Objective: In this review, we endeavor to do a literature search mainly focusing on keywords CD4+FOXP3+, CD4+CD127- , CD4+CD39+, CD4+LAP+ cancer immunotherapy, cancer immunotherapy strategies and articles published between 2015 to 2019 to add onto the minimal definition of human Treg, by an international workshop organized by collaborative immunoguiding program. Metho ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-8591.000028

Case Report

[Open Access](#) | [Case Report](#) | PTZAID:IJICR-6-126

[Invasive squamous cell carcinoma of the conjunctiva](#)

Published On: October 08, 2020 | Pages: 022 - 024

Author(s): Sonda Kammoun, Mona Rekik*, Saloua Ben Amor and Jamel Feki

Introduction: Squamous cell carcinoma of the conjunctiva is generally a low-grade tumor. Invasive squamous cell carcinoma, however, is uncommon, and intraocular extension has rarely been reported. We report a case of invasive squamous cell carcinoma of the conjunctiva with intra-ocular involvement in 94 year-old man. Methods: A case report.

Results: A 94-year-old ma ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-8591.000026

[Open Access](#) | [Case Report](#) | PTZAID:IJICR-6-125

[Lymphoepithelioma like Carcinoma of the Bladder – “A Case” Revisited](#)

Published On: September 14, 2020 | Pages: 019 - 021

Author(s): Mezhuneituo Raleng, Gaurav Aggarwal* and Sujoy Gupta

Lymphoepitheliomas are a type of undifferentiated carcinomas primarily described in the nasopharyngeal cavity. In the urinary tract, they are typically, but uncommonly seen in the urinary bladder, although occasional cases have been reported in the renal pelvis, ureter and urethra. Lymphoepithelioma-like carcinoma of the urinary bladder, first reported by Zuckerberg, ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-8591.000025

Short Communication

[Open Access](#) | [Short Communication](#) | PTZAID:IJICR-6-123

[New insights on assessing cancer risk of exposure to carcinogens from tobacco smoke inhalation](#)

Published On: June 03, 2020 | Pages: 014 - 015

Author(s): Samera Hussein Hamad*

Tobacco products is still the leading cause of deaths worldwide. 80% of lung cancer is attributed to tobacco smoking [1-3]. It has been found that the exposure to tobacco smoke is the primary etiologic factor responsible for lung cancer; and that the decline in the incidence and the mortality of lung cancer is associated with the decline in smoking [3,4]. At the same ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/2455-8591.000023

Open Access Mini Review PTZAID:IJICR-6-124

[Novel insights on use of doxorubicin to treat chemoresistant TNBC by Immunotherapy](#)

Published On: July 18, 2020 | Pages: 016 - 018

Author(s): Tze-Chen Hsieh* and Joseph M Wu

Doxorubicin, a WHO listed essential medicine, is used primarily as a combination of taxane/ anthracycline (doxorubicin)/cyclophosphamide (TAC), for treating malignant diseases including triple-negative breast cancer (TNBC). TNBC is a subtype of breast cancer categorized by deficient estrogen receptor, progesterone receptor, and epidermal growth factor receptor HER-2 [...

[Abstract View](#)

[Full Article View](#)

[DOI: 10.17352/2455-8591.000024](#)