

In this issue

Research Article

[Open Access](#) [Research Article](#) PTZAID:OJPM-3-116

[Clinical practice evaluation of opioids induced constipation management in patients with cancer: The EIO-50 project](#)

Published On: December 17, 2019 | Pages: 034 - 040

Author(s): Vicente Guillem, Francisco Javier Salvador Bofill, Margarita Feyjoo, Antonio Antón, Enrique Aranda, Carlos Camps, Alfredo Carrato, Manuel Constenla, Juan Jesús Cruz-Hernández, Eduardo Díaz-Rubio, Jesús García-Foncillas, Pere Gascón, Rafael López, Begoña Soler* and Yolanda Escobar

Introduction: Opioid-Induced Constipation (OIC) is a common distressing symptom associated with cancer pain treatment. Consensus guidelines for management of OIC are limited and little is known about the decision making of Health Care Professionals (HCP) for the management of OIC. The aim of this study was to learn about the diagnostic and treatment criteria of OIC ma ...

[Abstract View](#) [Full Article View](#) [DOI: 10.17352/ojpm.000016](#)

[Open Access](#) [Research Article](#) PTZAID:OJPM-3-115

[Analysis of care for cancer patients with opioid-induced constipation in clinical practice: EIO-PRAXIS Project](#)

Published On: December 06, 2019 | Pages: 027 - 033

Author(s): Enrique Aranda, Jesus Garcia-Foncillas, Antonio Anton, Carlos Camps, Alfredo Carrato, Manuel Constenla, Juan J Cruz-Hernandez, Eduardo Diaz-Rubio, Pere Gascon, Vicente Guillem, Rafael Lopez, Margarita Feyjoo, Begona Soler* and Javier Puente

Purpose: The objective of the study was to analyses how Opioid-Induced Constipation (OIC) is managed in cancer patients in clinical practice for comparison against the current recommendations for the management of patients with OIC.

...

[Abstract View](#) [Full Article View](#) [DOI: 10.17352/ojpm.000015](#)

[Open Access](#) [Research Article](#) PTZAID:OJPM-3-113

[Referred scapula pain from C6 or C7 cervical spinal stenosis](#)

Published On: September 04, 2019 | Pages: 021 - 023

Author(s): Marena Willeford* and Sierra Willeford

Objective: The aim of the present study was to determine if there exists referred medial scapula pain from C6 or C7 cervical spinal stenosis. Scapula pain of neurologic origin at present is felt to be mediated mainly by C5 through the dorsal scapular nerve. An interventional spine clinic noted a series of patients with medial scapula pain without C5 stenosis, however ...

[Abstract View](#) | [Full Article View](#) | DOI: [10.17352/ojpm.000013](https://doi.org/10.17352/ojpm.000013)

[Open Access](#) | [Research Article](#) | PTZAID:OJPM-3-110

[Observational study to analyse the opioid titration process in the treatment of breakthrough pain of cancer patients in clinical practice](#)

Published On: May 31, 2019 | Pages: 001 - 007

Author(s): Lara Iglesias-Docampo, Paola Pimentel-Cáceres, M^a Paola García-Coves, José Luis Firvida-Pérez, Julio José Lambea-Sorrosal, Antonio Andrés Rueda-Ramos, Iker López-Calderero, Juan Antonio Núñez-Sobrino, Antonio Javier Jiménez-López, Ana Cabezón-Álvarez and Begoña Soler-López*

Background: During the oncological disease, patients may suffer a special type of pain called breakthrough pain (BTP), defined as a transient exacerbation of pain that occurs either spontaneously, or in relation to a specific predictable or unpredictable trigger, despite relatively stable and adequately controlled background pain. Opioids are an essential part of BTP ...

[Abstract View](#) | [Full Article View](#) | DOI: [10.17352/ojpm.000010](https://doi.org/10.17352/ojpm.000010)

Review Article

[Open Access](#) | [Review Article](#) | PTZAID:OJPM-3-112

[Methods for saving opioids in the perioperative period via intravenous, via neuraxial and in nervous blocks](#)

Published On: September 03, 2019 | Pages: 015 - 020

Author(s): Borja Mugabure Bujedo*

The greatest challenge facing anesthesiologists in the current clinical context is to provide adequate perioperative care, achieving sufficient control of acute pain and at the same time make responsible use of intraoperative opioids. ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/ojpm.000012

[Open Access](#) | [Review Article](#) | PTZAID:OJPM-3-111

[Consensus statement on the management of breakthrough cancer pain: Assessment, treatment and monitoring recommendations](#)

Published On: August 26, 2019 | Pages: 008 - 014

Author(s): María Dolores López Alarcón*, Francisco Villegas Estevez, Vicente Domingo Triado, Pilar Blasco Segura, Genoveva Hernández Comes, Alfonso Berrocal Jaime, Ana Blasco Cordellat, Jorge Pastor Peidro, José López Torrecilla, Carlos Ferrer Albiach, Almudena Ruiz Sastre and Manuel Hernández Peris

Objective: Breakthrough cancer pain (BTcP) is still underestimated and a percentage of patients are not adequately treated, affecting their quality of life. The aim of this review was to generate a series of expert recommendations based on clinical experience and current scientific data on the management of BTcP. ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/ojpm.000011

Case Report

[Open Access](#) | [Case Report](#) | PTZAID:OJPM-3-114

[Malignant hyperthermia in young patient undergoing a surgical procedure under general anesthesia](#)

Published On: September 06, 2019 | Pages: 024 - 026

Author(s): Estéfano Arias Pastrana, Rosa Inés Carrascal Villero, Kevin Santiago Jiménez Escudero and Marlon Munera*

Malignant hyperthermia (MH) is a hypermetabolic disease induced by the administration of anesthetic drugs in patients with genetic susceptibility. Its incidence ranges from 0.02% to 0.0153%. Here, we report the case of an 18-year-old male patient with no previous relevant surgical, medical, or family history of disease, who while undergoing a scheduled orthopedic su ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/ojpm.000014