

In this issue

Research Article

[Open Access](#) [Research Article](#) PTZAID:OJT-5-137

**[“Perrotta Psychotherapeutic Protocol for Disorders of the Neurotic Area” \(PPP-DNA\): Proposal of protocol, profiles and clinical applications. Research](#)**

Published On: October 07, 2021 | Pages: 010 - 018

Author(s): Giulio Perrotta\*

Purpose: This research addresses the topic of anxiety, phobic and obsessive disorders. In this research, the theme is addressed to the psychopathological investigation of personalities, according to the PICI-2TA model (Perrotta Integrative Clinical Interviews, version 2-TA), the PAD-Q (Perrotta Affective Dependency Questionnaire), the PSM (Perrotta Sexual Matrix), the ...

[Abstract View](#) [Full Article View](#) [DOI: 10.17352/ojt.000037](#)

[Open Access](#) [Research Article](#) PTZAID:OJT-5-134

**[Traumas as social interactions](#)**

Published On: May 06, 2021 | Pages: 001 - 003

Author(s): Sam Vaknin\*

We react to serious mishaps, life altering setbacks, disasters, abuse, and death by going through the phases of grieving. Traumas are the complex outcomes of psychodynamic and biochemical processes. But the particulars of traumas depend heavily on the interaction between the victim and his social milieu. It would seem that while the victim progresses from denial to h ...

[Abstract View](#) [Full Article View](#) [DOI: 10.17352/ojt.000034](#)

Review Article

[Open Access](#) [Review Article](#) PTZAID:OJT-5-138

## [Splenic trauma: Definition, classifications, clinical profiles and best treatments](#)

Published On: October 21, 2021 | Pages: 019 - 036

Author(s): Giulio Perrotta\*, Emanuele Guerrieri and Mario Guerrieri

The spleen is an organ commonly injured in abdominal trauma of the upper left quadrant and until just under two decades the first choice was always splenectomy; however, based on new research and clinical experience, there is a tendency to preserve the spleen as much as possible, precisely because of its immune function and risk of infection. On the basis of the traum ...

[Abstract View](#) | [Full Article View](#) | DOI: [10.17352/ojt.000038](https://doi.org/10.17352/ojt.000038)

### Case Report

[Open Access](#) [Case Report](#) PTZAID:OJT-5-136

## [Pseudoaneurism of the brachial artery mimicking a soft tissue sarcoma: Case report](#)

Published On: September 14, 2021 | Pages: 006 - 009

Author(s): John Musubao Katsuva\*, Simplicie Kighoma Vuhaka and Sosthene Tsongo Vululi

The current case is one of the rare clinical presentations of the brachial artery pseudo aneurysm presentation in children Pseudo-aneurism is one of the late complications of a missed or untreated arterial injury. The diagnosis is suggested in the presence of clinical signs such as: an expending, ill-defined mass with or without pulsation; bruit, pain, paresthesia, or ...

[Abstract View](#) | [Full Article View](#) | DOI: [10.17352/ojt.000036](https://doi.org/10.17352/ojt.000036)

[Open Access](#) [Case Report](#) PTZAID:OJT-5-135

## [May a dissociative disorder contain sexual abuse of children through internet? A case report](#)

Published On: July 29, 2021 | Pages: 004 - 005

Author(s): John E Berg\*

Post-traumatic stress after abuse early in life, may be forgotten until late in life. Dissociative actions may be observed. An example of this is demonstrated in a case report from a resident psychiatric facility in Northern Norway. Under Norwegian

Law, acts under dissociation is not regarded as being not fit for standing trial, but may be deemed a partial excuse. In ...

[Abstract View](#)

[Full Article View](#)

[DOI: 10.17352/ojt.000035](#)